



Aberystwyth University

Comment on "Liquid-Liquid Phase Transition in Supercooled Yttria-Alumina"

Greaves, G. N.; Wilding, M. C.; Hennes, L.; Langstaff, D.; Kargl, F.; Benmore, C. J.; Weber, J. K. R.

Published in:
Physical Review Letters

DOI:
[10.1103/PhysRevLett.106.119601](https://doi.org/10.1103/PhysRevLett.106.119601)

Publication date:
2011

Citation for published version (APA):

Greaves, G. N., Wilding, M. C., Hennes, L., Langstaff, D., Kargl, F., Benmore, C. J., & Weber, J. K. R. (2011). Comment on "Liquid-Liquid Phase Transition in Supercooled Yttria-Alumina". *Physical Review Letters*, 106(11), [119601]. <https://doi.org/10.1103/PhysRevLett.106.119601>

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Comment on “Liquid-Liquid Phase Transition in Supercooled Yttria-Alumina”

We welcome efforts made by Barnes *et al.* to study liquid-liquid transitions (LLT) in supercooled Y_2O_3 - Al_2O_3 (100- x) or AY_x melts reported in a recent Letter [1]. The first order LLT in AY_{20} , which we have identified *in situ* under equilibrium conditions [2,3] has a well-defined temperature T_{LL} at ambient pressure and a critical point at negative pressure. The LLT is also composition dependent, rising as x falls [3,4]. We identified this at 1788 K from (i) a peak in the small angle x-ray scattering (SAXS) intensity, (ii) a discontinuity in the structure factor $S(Q)$, and (iii) a “polyamorphic rotor” caused by periodic LLTs [Fig. 1(a)]—the flipping time and associated temperature spikes yielding the LLT density and entropy discontinuities in agreement with *ex situ* experiments [4]. Barnes *et al.* used similar experiments but could not reproduce (i) or (ii) in AY_{20} at 1788 K and attributed (iii) to 60 K instabilities sometimes encountered during sample conditioning [1]. We consider their null results for AY_{20} are due to (A) large neutron beam sizes in small angle neutron scattering (SANS), and (B) doubts in AY_x composition. Moreover, comparison with our work [2] is obscured in [1] by reliance on apparent temperatures uncorrected for emissivity [5] and by inconsistent molar normalization leading to flaws in modelling LLTs from our data [3].

(A) In their SANS measurements [1] the mm radius droplet was overspillied by a 2 mm radius 4.5 \AA beam. Because of the spherical liquid surface this results in total external reflection and cross fire contamination up to at least $2 \times 0.013 = 0.026 \text{ \AA}^{-1}$. Sub mm focused SAXS has none of these disadvantages [2] which is why it is sensitive to the rise and fall in scatter that occurs below 0.03 \AA^{-1} [2] at the LLT (Fig. 1). Importantly, polyamorphic rotor action was recorded by L. Hennet during the SANS experiments with periodic 150 K spikes centered at 1940 K [Fig. 1(b)]. Polyamorphic rotors have large repetitive spikes thermally distinct from the oscillations illustrated in [1] which sometimes occur when molten drops contain inclusions.

(B) The sample preparation method of fusing 85 mg drops from weight-matched beads of separate oxides [6] is unreliable without validation of recovered samples. We fused mm radius drops by weighing material from 5 g powdered batches following repeated sintering and re-grinding. Composition checks post experiment confirm $x < \text{mol } 1\%$ accuracy. $S(Q)$'s of AY_x melts at 2300 K are composition sensitive (Fig. 1). Measurements at 11-ID-C (APS) [2] and at ID11 (ESRF) [7] are in good agreement, with the positions of the first and second peaks Q_1 and Q_2 , and size of the principal peak $S(Q_1)$ all scaling almost linearly with x , except for values fitted from the peak maxima at 2300 K from [1] Fig. 1. Rather than AY_{20} , these triangulate with the composition $AY_{14} \pm 1$. We already investigated AY_{15} until crystallization intervened at 1927 K [2] close to the rotor temperature shown in




FIG. 1 (color online). Left. Upper: SAXS for AY_{20} ■ and AY_{15} △ from [2] scaled to SANS for AY_{20} ● from [1], together with LLT rotor motion (A and B). Lower: $S(Q)$ for AY_x liquids from [2] corrected for detector scattering angle thickness, with compositional changes marked. Right: trends in Q_2 , Q_1 and $S(Q_1)$ with x . 2σ errors for Q_1 and Q_2 from peak-fitting and for $S(Q_1)$ from [1] within symbol size. Arrows between closed and open symbols \updownarrow show shifts in $S(Q_1)$ for alternative compositions $x = 15 \leftrightarrow 20$. Horizontal arrows show $x = 14\% \pm 1\%$ as the likely composition for the Letter [1] rather than AY_{20} .

Fig. 1(b). If 1940 K marks the LLT for AY_{15} then, with our observation of 1788 K for AY_{20} [2], these results further demonstrate that for AY_x liquids T_{LL} rises as x falls [3,4], and provide scope for future collaboration.

G. N. Greaves,^{1,2} M. C. Wilding,² L. Hennet,³
D. Langstaff,² F. Kargl,² C. J. Benmore,⁴ and
J. K. R. Weber^{4,5}

¹Department Materials Science & Metallurgy, University of Cambridge, Cambridge CB2 3QZ, United Kingdom

²CAFMaD, Aberystwyth University, Aberystwyth SY23 3BZ, United Kingdom

³CNRS-CEMHTI and University of Orléans, 45071 Orléans, France

⁴Argonne National Laboratory, Argonne, Illinois 60439, USA

⁵Materials Development, Inc., Arlington Heights, Illinois 60004, USA

Received 12 December 2009; revised manuscript received

1 June 2010; published 18 March 2011

DOI: 10.1103/PhysRevLett.106.119601

PACS numbers: 64.70.Ja, 61.05.cp, 61.20.Lc, 64.60.My

[1] A. C. Barnes *et al.*, *Phys. Rev. Lett.* **103**, 225702 (2009).

[2] G. N. Greaves *et al.*, *Science* **322**, 566 (2008).

[3] G. N. Greaves *et al.*, *J. Non-Cryst. Solids* **357**, 435 (2011).

[4] S. Aasland and P. F. McMillan, *Nature (London)* **369**, 633 (1994).

[5] Emissivity = 0.92 for AY_x melts: J. K. Weber *et al.*, *J. Am. Ceram. Soc.* **83**, 1868 (2004).

[6] L. B. Skinner *et al.*, *J. Phys. Condens. Matter* **20**, 205103 (2008).

[7] V. Cristiglio, Ph.D. thesis, Université d’Orléans, 2008.